

THE DUAL GEOMETRY OF BOOLEAN SEMIRINGS

DANIEL J. CLOUSE AND FERNANDO GUZMÁN

ABSTRACT. It is well known that the variety of Boolean semirings, which is generated by the three element semiring $\mathbb{S}$, is dual to the category of partially Stone spaces. We place this duality in the context of natural dualities. We begin by introducing a topological structure $\underline{\mathbf{S}}$ and obtain an optimal natural duality between the quasi-variety $ISP(\mathbb{S})$ and the category $IS_cP^+(\underline{\mathbf{S}})$. Then we construct an optimal and very small structure $\underline{\mathbf{S}}_{os}$ that yields a strong duality. The geometry of some of the partially Stone spaces that take part in these dualities is presented, and we call them "hairy cubes", as they are n -dimensional cubes with unique incomparable covers for each element of the cube. We also obtain a polynomial representation for the elements of the hairy cube.

1. INTRODUCTION

Extensions of the concept of a *Boolean Ring* to include semirings has been done in several different directions. One source of diversity are the different definitions of *semiring*. The other is how they get connected to Boolean rings. We will use the concept of semiring commonly used in formal languages and automata theory, that is, the only thing missing in order to be a ring is the existence of additive inverses (see [3] and [6]). As in Guzman [4], we will denote by $\mathcal{BSR}$ the variety generated by the two 2-element semirings, and will call it the variety of Boolean semirings. It turns out that this variety is also generated by a 3-element semiring with carrying set $S = \{0, h, 1\}$, that we denote $\mathbb{S}$. The semiring $\mathbb{S}$ will play a crucial role in this paper.

In [4], following the ideas of Stone [7] in his now famous "*Stone representation theorem*", a duality is established between the category $\mathcal{BSR}$ of Boolean semirings and the category $\mathcal{PSS}$ of partially Stone spaces. On the other hand, Clark and Davey [1] present a thorough study of natural dualities between algebraic and topological quasi-varieties. It is the goal of this paper to place the duality from [4] in the much richer context of [1].

A *structured topological space* consists of

$$\mathbf{X} = \langle X; G, H, R, T \rangle$$

where $\langle X, T \rangle$ is a topological space, G is a set of finitary (total) operations on X , H is a set of finitary partial operations on X and R is a set of finitary relations on X . The arities of the operations, partial operations, and relations define the type of $\mathbf{X}$. Given a finite discrete structured topological space $\mathbf{X}$, we denote by $IS_cP^+(\mathbf{X})$ the category of closed substructures of non-empty products of copies of $\mathbf{X}$.

Date: January 28, 2008.

2000 Mathematics Subject Classification. Primary:06D50, 08C15; Secondary:08C05, 18A40, 06E15.

Key words and phrases. Boolean Semiring, Duality, Strong duality, Optimal duality.

On one side of the duality we will have the quasi-variety $\mathcal{A} = ISP(\mathbb{S})$ generated by $\mathbb{S}$. On the other side, we will have the category of structured topological spaces $\mathcal{X} = IS_cP^+(\underline{\mathbb{S}})$, generated by some appropriate structure $\underline{\mathbb{S}} = \langle S; G, H, R, T \rangle$ having $S = \{0, h, 1\}$ as underlying set and T the discrete topology.

We begin by naming three binary relations on S : r_1 , r_2 , and r_3 , and prove the following result.

Theorem 1. *The structure $\underline{\mathbb{S}} = \langle S; \{r_1, r_2, r_3\}, T \rangle$ yields an optimal natural duality on $\mathcal{A}$.*

The proof is in Section 2. The duality is optimal in the sense that if any one of the relations were to be deleted from the structure of $\underline{\mathbb{S}}$, duality would be lost.

In the duality $\mathcal{BSR} \rightleftharpoons \mathcal{PSS}$, a Boolean semiring A is mapped to the set of prime filters of A (recall that a Boolean semiring can be viewed as a partially complemented distributive lattice. See [4]). For finite A , every prime filter is the upset of a join-irreducible element. When $\mathbf{X}$ is a closed substructure of a finite power of $\underline{\mathbb{S}}$, we would like to describe the join-irreducible elements of $\mathcal{X}(\mathbf{X}, \underline{\mathbb{S}})$. We denote the set of all of them by $\mathcal{X}(\mathbf{X}, \underline{\mathbb{S}})_J$. In **Theorem 2**, join-irreducible elements of $\mathcal{X}(\underline{\mathbb{S}}^n, \underline{\mathbb{S}})$ is given; see Section 2. In [2] a description of $\mathcal{X}(\mathbf{X}, \underline{\mathbb{S}})_J$ is given for any $\mathbf{X} \in \mathcal{X}$ that is a closed substructure of a finite power of $\underline{\mathbb{S}}$. This will appear in a subsequent paper. Here we lay the foundation for those results, a description of the meet semilattice $\mathcal{X}(\underline{\mathbb{S}}^n, \underline{\mathbb{S}})_J$, that we call the “hairy cube”. It consists of an n -dimensional cube covered by “hairs”. More precisely,

Theorem 3. *The poset $\mathcal{X}(\underline{\mathbb{S}}^n, \underline{\mathbb{S}})_J$ consists of two parts: the “base” Y^n which is an n -cube and the “hairs” $\mathcal{X}(\underline{\mathbb{S}}^n, \underline{\mathbb{S}})_J \setminus Y^n$, which are pairwise incomparable. Each element of the base is covered by a unique hair. Each hair covers a unique base element.*

In **Theorem 4** it is shown that these partial order properties of $\mathcal{X}(\underline{\mathbb{S}}^n, \underline{\mathbb{S}})_J$ completely determine it as a partially Stone space. We close Section 3 with a polynomial representation of the join-irreducible elements of $\mathcal{X}(\underline{\mathbb{S}}^n, \underline{\mathbb{S}})$. See **Theorem 5**.

In Section 4 we first establish that the duality in Theorem 1 is neither a full nor a strong duality. Then we discuss why this is true and how that duality can easily be upgraded to a strong duality, following some of the ideas of [1]. Then we show how to construct an optimal and very small structure $\underline{\mathbb{S}}_{os}$ that yields a strong duality on $\mathcal{A}$. Despite the fact that $\mathbb{S}$ is not subdirectly irreducible, $\text{Irr}(\mathbb{S})$ is 2, $\underline{\mathbb{S}}_{os}$ consists of a single relation r_2 , and a single partial operation λ_1 .

Theorem 6. *Let T denote the discrete topology and $\underline{\mathbb{S}}_{os} = \langle S; \{r_2\}, \{\lambda_1\}, T \rangle$. Then $\underline{\mathbb{S}}_{os}$ yields an optimal strong duality on $\mathcal{A}$.*

Finally, we discuss why the “Hairy Cube” will persist in that strong duality.

Corollary 6. *Let $\mathcal{X}_{os} = IS_cP^+(\underline{\mathbb{S}}_{os})$, then for any $n \in \mathbb{N}$, $\mathcal{X}_{os}(\underline{\mathbb{S}}^n, \underline{\mathbb{S}})$ is the n -dimensional Hairy Cube.*

The results in this paper and in [2] greatly expand our understanding of the dual equivalence between the variety of Boolean Semirings and the category of Partially Stone Spaces

established in [4]. They also complete a large initial step for investigating the strong duality we establish between $\mathcal{A}$ and $\mathcal{X}_{os}$.

1.1. Notation.

Since some of the arguments in the paper are of an inductive nature, we need a convenient notation to move back and forth between functions $S^{n-1} \rightarrow S$ and functions $S^n \rightarrow S$.

Given $n \in \mathbb{N}$, $\Phi : S^n \rightarrow S$, $a \in S$, and $x \in S^{n-1}$ we denote by Φ_a the map

$$\begin{aligned} \Phi_a : \quad S^{n-1} &\rightarrow S \\ (x_1, \dots, x_{n-1}) &\mapsto \Phi(a, x_1, \dots, x_{n-1}) \end{aligned}$$

and by Φ_x the map

$$\begin{aligned} \Phi_x : \quad S &\rightarrow S \\ a &\mapsto \Phi(a, x_1, \dots, x_{n-1}) \end{aligned}$$

Given $\psi, \psi', \psi'' : S^{n-1} \rightarrow S$, we denote by (ψ, ψ', ψ'') the map

$$\begin{aligned} (\psi, \psi', \psi'') : \quad S^n &\rightarrow S \\ (x_1, x_2, \dots, x_n) &\mapsto \begin{cases} \psi(x_2, \dots, x_n) & \text{if } x_1 = 0 \\ \psi'(x_2, \dots, x_n) & \text{if } x_1 = h \\ \psi''(x_2, \dots, x_n) & \text{if } x_1 = 1 \end{cases} \end{aligned}$$

Note that for any $\Phi : S^n \rightarrow S$ we have $\Phi = (\Phi_0, \Phi_h, \Phi_1)$. In particular, when $n = 1$ we may write any $\Phi : S \rightarrow S$ as a triple of elements of S ; see, for example, Lemma 3. Since $\Phi_a(x) = \Phi_x(a)$ for any $x \in S^{n-1}$ and $a \in S$, we have $\Phi_x = (\Phi_0(x), \Phi_h(x), \Phi_1(x))$. When $\psi : S^{n-1} \rightarrow S$ is a term function of $\mathbb{S}$ on $(n-1)$ variables, then $\Psi = (\psi, \psi, \psi)$ is the same term function viewed as a term function on n variables (with the first one absent). We call Ψ the n -ary version of ψ .

For $1 \leq i \leq n \in \mathbb{N}$ we denote by Π_i^n the i -th projection map

$$\begin{aligned} \Pi_i^n : \quad S^n &\rightarrow S \\ (x_1, \dots, x_n) &\mapsto x_i \end{aligned}$$

For any binary relation $r \subseteq S^2$, we denote by r^{-1} the inverse relation $\{(y, x) | (x, y) \in r\}$.

Given a structured topological space $\mathbf{X}$, and $\mathbf{Y} \in IS_c P^+(\mathbf{X})$, for any operation, partial operation or relation λ of the structure $\mathbf{X}$, we denote by $\lambda^{\mathbf{Y}}$ the corresponding operation, partial operation or relation on $\mathbf{Y}$.

Definition 1. [4] *A partially complemented distributive lattice is a type $\langle 0, 0, 0, 2, 2 \rangle$ algebra $A = \langle A; 0, h, 1, \vee, \wedge \rangle$ such that $\langle A; 0, 1, \vee, \wedge \rangle$ is a bounded distributive lattice and $\langle [h, 1]; h, 1, \vee, \wedge \rangle$ is a complemented distributive lattice, i.e., a Boolean algebra where $[h, 1] = \{a \in A | h \leq a \leq 1\}$.*

We also wish to note here that it is shown in [4] that

In any partially complemented distributive lattice one can define a unary (bar) operation in terms of the complement operation $'$ in $[h, 1]$:

$$\overline{x} = (x \vee h)'$$

It satisfies two useful identities can be defined as in the following:

Lemma 1. [4] *Given a partially complemented distributive lattice $\langle A; 0, h, 1, \vee, \wedge \rangle$, the bar operation satisfies*

L1) $x \vee \overline{x} = 1$, and

L2) $x \wedge \overline{x} = x \wedge \overline{1}$.

These two properties characterize partially complemented distributive lattices. From these results it is derived that $\mathcal{BSR}$ is dually equivalent to the category of Partially Stone Spaces, $\mathcal{PSS}$, and we denote this dual equivalence as $\mathcal{BSR} \rightleftharpoons \mathcal{PSS}$.

In $\mathcal{BSR} \rightleftharpoons \mathcal{PSS}$, the functor from $\mathcal{BSR}$ to $\mathcal{PSS}$ takes any partially complemented distributive lattice A and maps it to $\text{pt}(\text{Idl}(A))$, the set of prime filters of A . These prime filters are difficult to characterize for arbitrary powers of $\mathbb{S}$. For example, if the cardinality of the indexing set is at least countably infinite, every prime filter has an infinite descending chain and the existence of elements of finite support is unclear. Hence our current understanding of $\mathcal{BSR}$ inherent in this representation is not entirely satisfactory.

1.2. Natural Dualities.

In this paper we will follow very closely the ideas of [1] for constructing natural dualities. The basic idea is to impose on the carrier S of the semiring $\mathbb{S}$, the discrete topology together with operations, partial operations and relations to form a dual topological structure $\underline{\mathbb{S}}$ as the generator of the dual category $\mathcal{X}$. More specifically, $\mathcal{X} = IS_cP^+(\underline{\mathbb{S}})$ is the category of isomorphic copies, topologically closed substructures of non-empty products of copies of $\underline{\mathbb{S}}$. Following this construction, we will have a dual adjunction $\langle D, E, e, \epsilon \rangle$ between the categories $\mathcal{A}$ and $\mathcal{X}$ with the many desirable properties [1, 1.5.3]. One further property we desire is that for any $A \in \mathcal{A}$, A is isomorphic to $ED(A) = \mathcal{X}(\mathcal{A}(A, \mathbb{S}), \underline{\mathbb{S}})$. If the dual adjunction $\langle D, E, e, \epsilon \rangle$ satisfies this property, it is called a dual representation of $\mathcal{A}$ in $\mathcal{X}$. In this case we say that $\underline{\mathbb{S}}$ yields a (natural) duality on $\mathcal{A}$. If it is also true that for any $X \in \mathcal{X}$, X is isomorphic to $DE(X) = \mathcal{A}((X, \underline{\mathbb{S}}), \mathbb{S})$, we say that $\underline{\mathbb{S}}$ yields a full duality on $\mathcal{A}$. Thus $\underline{\mathbb{S}}$ yields a full duality on $\mathcal{A}$ if it yields a duality on $\mathcal{A}$ which is a dual equivalence. Finally, if $\underline{\mathbb{S}}$ yields a full duality on $\mathcal{A}$ and it is injective in $\mathcal{X}$, $\underline{\mathbb{S}}$ is said to yield a strong duality on $\mathcal{A}$.

We will construct three dualities, each one coming from a different topological structure. In all three of them the algebra side of the duality will be $\mathcal{A} = ISP(\mathbb{S})$. The first topological structure, $\underline{\mathbb{S}}$, yields an optimal (natural) duality $\mathcal{X} = IS_cP^+(\underline{\mathbb{S}}) \rightleftharpoons \mathcal{A}$. The second one, $\underline{\mathbb{S}}_s$, yields a strong duality $\mathcal{X}_s = IS_cP^+(\underline{\mathbb{S}}_s) \rightleftharpoons \mathcal{A}$. The third one, $\underline{\mathbb{S}}_{os}$, yields an optimal strong duality $\mathcal{X}_{os} = IS_cP^+(\underline{\mathbb{S}}_{os}) \rightleftharpoons \mathcal{A}$. In the first duality, labeling the appropriate contravariant functors D and E , A is isomorphic to $ED(A) = \mathcal{X}(\mathcal{A}(A, \mathbb{S}), \underline{\mathbb{S}})$. The situation is displayed in the diagram below.

$$\mathcal{X} \xrightarrow{D, E} \mathcal{A} = ISP(\mathbb{S}) \hookrightarrow HSP(\mathbb{S}) = \mathcal{BSR} \rightleftharpoons \mathcal{PSS}$$

Similar remarks hold for the other two dualities. Moreover, in Corollary 6 we show that $\mathcal{X}_{os}(\underline{\mathbb{S}}_{os}^n, \underline{\mathbb{S}}) = \mathcal{X}(\underline{\mathbb{S}}^n, \underline{\mathbb{S}})$.

Recall that for any $\mathbf{X} \in IS_cP^+(\underline{\mathbb{S}})$, $E(\mathbf{X}) = \mathcal{X}(\mathbf{X}, \underline{\mathbb{S}}) \leq \mathbb{S}^{\mathbf{X}}$, and $DE(\mathbf{X})$ is the set of prime filters of $\mathcal{X}(\mathbf{X}, \underline{\mathbb{S}})$. As a result, we desire that the structure placed on $\underline{\mathbb{S}}$ will be sufficient so that we can characterize the prime filters of $\mathcal{X}(\mathbf{X}, \underline{\mathbb{S}})$, and thereby trace their images and the

image of $\mathcal{A}$ in $\mathcal{PSS}$. In [4] it is shown that for a finite partially complemented distributive lattice $\mathbb{L}$, the partially Stone space $[X, Y]$ corresponding to $\mathbb{L}$ under the $\mathcal{BSR} \rightleftharpoons \mathcal{PSS}$ duality, has $X = \mathbb{L}_J$ and $Y = \{x \in X | x \leq h\}$. The topology of this space is $\mathcal{T} = \{\phi(I) | I \in \text{Idl}(\mathbb{L})\}$ where $\phi(I) = \{p \in X | p \cap I \neq \emptyset\}$ for any $I \in \text{Idl}(\mathbb{L})$. We call this topology the Stone Space topology. In particular, when $\mathbb{L} = \mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})$, we identify the prime filters of $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})$ with its join-irreducible elements. A major portion of this paper is devoted to characterizing the join-irreducible elements of $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})$; we denote the set of such elements by $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$.

2. ESTABLISHING THE DUALITY AND SOME FACTS ABOUT MORPHISMS

First of all, we note that $t(x, y, z) = xy + yz + xz$ is a ternary near-unanimity term on $\mathbb{S}$. This property will allow us to use many of the results in [1] in the construction of the dual representations that we seek. We further note that this property implies that $\mathcal{BSR}$ is a congruence distributive variety and finite products of $\mathbb{S}$ are skew-free. Arbitrary products of $\mathbb{S}$ are known to the authors not to be skew-free, but the counterexample is outside of the scope of this paper.

We now define the first structure $\underline{\mathbf{S}}$ that we will show yields a duality on $\mathcal{A}$. We will also show that this duality is optimal, in the sense that if any single relation were to be deleted from $\underline{\mathbf{S}}$, duality would be lost.

Definition 2. Define the following subsets of $\mathbb{S}^2$:


$$\begin{aligned} r_1 &= \mathbb{S}^2 - \{(1, 0)\}; \\ r_2 &= \mathbb{S}^2 - \{(h, 0), (1, 0)\}; \\ r_3 &= \mathbb{S}^2 - \{(0, 1), (h, 1), (1, 0), (1, h)\}. \end{aligned}$$

Let $\underline{\mathbf{S}} = \langle S; \{r_1, r_2, r_3\}, \mathcal{T} \rangle$ where $\mathcal{T}$ is the discrete topology.

The following result can be shown by straightforward counting and closure calculations.

Lemma 2. The following is the lattice of subalgebras of $\mathbb{S}^2$:

The Lattice of Subalgebras of $\mathbb{S}^2$



Consider now the lattice of subalgebras of $\mathbb{S}^2$ coupled with the following facts:

- if a morphism preserves a binary relation r then it also preserves r^{-1} , and
- if a morphism preserves two k -ary relations r and s then it also preserves their intersection $r \cap s$.


From this we see that if we use only the set of binary relations r_1 , r_2 , and r_3 from Definition 2 as structure, then the structure $\underline{\mathbf{S}}$ will entail all subalgebras of $\mathbb{S}^2$. Hence by the M-Shift Duality Lemma [1, 2.4.2] and the NU Duality Theorem [1, 2.3.4] we get the following proposition:

Proposition 1. $\underline{\mathbf{S}}$ yields a duality on $\mathcal{A}$ and $\underline{\mathbf{S}}$ is injective in $\mathcal{X}$.

The next lemma determines the elements of $\mathcal{X}(\underline{\mathbf{S}}, \underline{\mathbf{S}}) \leq \mathbb{S}^S$. The proof is a simple verification of which functions $f : S \rightarrow S$ preserves the relations of $\underline{\mathbf{S}}$.

Lemma 3. *The following is the lattice $\mathcal{X}(\underline{\mathbf{S}}, \underline{\mathbf{S}})$:*

The Lattice $\mathcal{X}(\underline{\mathbf{S}}, \underline{\mathbf{S}})$



Before we investigate the lattice $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$, we want to show that the duality yielded by $\underline{\mathbf{S}}$ on $\mathcal{A}$ is optimal. From the Preduality Theorem [1, 1.5.2] and the First Duality Theorem [1, 2.2.2] we get the following lemma.

Lemma 4. *If $\underline{\mathbf{S}}'$ is a structure that yields duality on $\mathcal{A}$, then the finitary term functions on $\underline{\mathbf{S}}$ must be exactly the morphisms from finite powers of $\underline{\mathbf{S}}'$ into $\underline{\mathbf{S}}'$. Therefore, for any $n \in \mathbb{N}$*

$$\mathcal{X}'(\underline{\mathbf{S}}^n, \underline{\mathbf{S}}') = \mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}}).$$

Theorem 1. *The structure $\underline{\mathbf{S}} = \langle S; \{r_1, r_2, r_3\}, \mathcal{T} \rangle$ yields an optimal natural duality on $\mathcal{A}$.*

Proof. Using Lemmas 3 and 4 we can show that if we eliminate r_2 or r_3 from $\underline{\mathbf{S}}$ to form $\underline{\mathbf{S}}'$, duality will be lost. The map $(1, h, 1)$ preserves r_1 and r_2 , but not r_3 . The map $(0, 0, h)$ preserves r_1 and r_3 , but not r_2 .

Now let $\underline{\mathbf{S}}' = \langle S; \{r_2, r_3\}, \mathcal{T} \rangle$, $\mathcal{X}' = IS_e P^+(\underline{\mathbf{S}}')$ and suppose that $\underline{\mathbf{S}}'$ yields a duality on $\mathcal{A}$. By the Duality and Entailment Theorem [1, 2.4.3], $\{r_2, r_3\}$ must entail r_1 . By the

Entailment Lemma [1, 2.4.4], for any $X \subseteq S^2$ and $\alpha \in \mathcal{X}'(X, \underline{\mathbf{S}}')$, α must preserve r_1 . Let $X = \{(h, 0), (0, 1)\}$ and define $\alpha : X \rightarrow \underline{\mathbf{S}}$ by $\alpha((h, 0)) = 1$ and $\alpha((0, 1)) = 0$. Then α preserves r_2 and r_3 , but not r_1 . Therefore, we cannot retain duality without r_1 . $\square$

Upon viewing Lemma 3, the Preduality Theorem and the First Duality Theorem, noting that h is a nullary operation of $\mathbf{S}$ and recalling that relations are defined pointwise, it is easy to see that Lemma 5 holds. This lemma gives some recursive information about $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})$, just enough for our needs.

Lemma 5. *Let $n > 1$.*


- (1) *Given $\Phi \in \mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})$ and $x \in S^{n-1}$, we have $\Phi_x \in \mathcal{X}(\underline{\mathbf{S}}, \underline{\mathbf{S}})$.*
- (2) *Given $\Phi \in \mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})$, and $a \in S$, we have $\Phi_a \in \mathcal{X}(\underline{\mathbf{S}}^{n-1}, \underline{\mathbf{S}})$, and*
 - (a) $\Phi_0 \vee (\Phi_1 \wedge h) = \Phi_h$, *hence* $\Phi_0 \leq \Phi_h$.
 - (b) $\Phi_0 \wedge h \leq \Phi_1$.

Moreover,

 - (c) *If $\Phi \leq h$ then $\Phi_h = \Phi_1$.*
 - (d) *If $\Phi \not\leq h$ then for every $x \in S^{n-1}$ either $\Phi_0(x) \leq \Phi_h(x) \leq \Phi_1(x)$ or $\Phi_0(x) = \Phi_h(x) = 1$ and $\Phi_1(x) = h$.*
- (3) *Given $\psi \in \mathcal{X}(\underline{\mathbf{S}}^{n-1}, \underline{\mathbf{S}})$, let $\Psi = (\psi, \psi, \psi)$ be the n -ary version of ψ . We have that $\Psi \wedge h \wedge \Pi_1, \Psi \wedge h \wedge \overline{\Pi_1}, \Psi \wedge \Pi_1, \Psi \wedge \overline{\Pi_1} \in \mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})$, and*
 - (a) $\Psi \wedge h \wedge \Pi_1 = (0, \psi \wedge h, \psi \wedge h)$
 - (b) $\Psi \wedge h \wedge \overline{\Pi_1} = (\psi \wedge h, \psi \wedge h, \psi \wedge h)$
 - (c) $\Psi \wedge \Pi_1 = (0, \psi \wedge h, \psi)$
 - (d) $\Psi \wedge \overline{\Pi_1} = (\psi, \psi, \psi \wedge h)$

We now have enough results to begin our work characterizing $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$. We begin with a simple but useful Corollary to Lemma 3.

Corollary 1. *The following diagram is the poset $\mathcal{X}(\underline{\mathbf{S}}, \underline{\mathbf{S}})_J$:*



Proposition 2. *Let $n > 1$.*

- (1) *If $\psi \in \mathcal{X}(\underline{\mathbf{S}}^{n-1}, \underline{\mathbf{S}})_J$ and $\Psi = (\psi, \psi, \psi)$ is the n -ary version of ψ , then $\Psi \wedge \Pi_1 = (0, \psi \wedge h, \psi)$ and $\Psi \wedge \overline{\Pi_1} = (\psi, \psi, \psi \wedge h)$ are in $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$.*

- (2) If $\Phi \in \mathcal{X}(\mathbf{S}^n, \mathbf{S})_J$ with $\Phi \leq h$, then there is $\psi \in \mathcal{X}(\mathbf{S}^{n-1}, \mathbf{S})_J$ with $\psi \leq h$ such that $\Phi = (0, \psi, \psi) = \Psi \wedge \Pi_1$ or $\Phi = (\psi, \psi, \psi) = \Psi \wedge \overline{\Pi_1}$, where $\Psi = (\psi, \psi, \psi)$ is the n -ary version of ψ .
- (3) If $\Phi \in \mathcal{X}(\mathbf{S}^n, \mathbf{S})_J$ with $\Phi \not\leq h$, then there is $\psi \in \mathcal{X}(\mathbf{S}^{n-1}, \mathbf{S})_J$ with $\psi \not\leq h$ such that $\Phi = (0, \psi \wedge h, \psi) = \Psi \wedge \Pi_1$ or $\Phi = (\psi, \psi, \psi \wedge h) = \Psi \wedge \overline{\Pi_1}$, where $\Psi = (\psi, \psi, \psi)$ is the n -ary version of ψ .

Proof. 1. Assume $\psi \in \mathcal{X}(\mathbf{S}^{n-1}, \mathbf{S})$ is join-irreducible.

Consider first the case $\Phi = (0, \psi \wedge h, \psi)$, and suppose $\Phi = \Gamma \vee \Delta$ with $\Gamma, \Delta \in \mathcal{X}(\mathbf{S}^n, \mathbf{S})$ and $\Gamma, \Delta < \Phi$. We have

$$\Gamma_0 = \Delta_0 = 0, \quad \Gamma_h \vee \Delta_h = \psi \wedge h, \quad \Gamma_1 \vee \Delta_1 = \psi.$$

Without loss of generality, we have $\Gamma_1 = \psi$ and therefore $\Gamma_h < \psi \wedge h$, i.e. there is $x \in S^{n-1}$ such that

$$0 \leq \Gamma_h(x) < \psi(x) \wedge h \leq \psi(x), h$$

So, we must have $\Gamma_h(x) = 0$. By Lemma 5.1, $\Gamma_x \in \mathcal{X}(\mathbf{S}, \mathbf{S})$ and by Lemma 3 $\Gamma_x = (0, 0, 0)$ which gives $0 = \Gamma_1(x) = \psi(x)$, a contradiction.

The case $\Phi = (\psi, \psi, \psi \wedge h)$ can be handled in the same way, except that instead of going from Γ_1 to Γ_h , one goes from Γ_0 to Γ_1 .

2. Assume $\Phi \in \mathcal{X}(\mathbf{S}^n, \mathbf{S})$ is join-irreducible, with $\Phi \leq h$.

By Lemma 5.2 $\Phi_h = \Phi_1 \in \mathcal{X}(\mathbf{S}^{n-1}, \mathbf{S})$, call it ψ , and $0 \leq \Phi_0 \leq \psi$. Moreover, $\psi \leq h$. By Lemma 5.3 we have (Φ_0, Φ_0, Φ_0) and $(0, \psi, \psi)$ are in $\mathcal{X}(\mathbf{S}^n, \mathbf{S})$, and

$$(\Phi_0, \Phi_0, \Phi_0) \vee (0, \psi, \psi) = (\Phi_0, \psi, \psi) = \Phi$$

Therefore, either $\Phi = (0, \psi, \psi)$ or $\Phi = (\Phi_0, \Phi_0, \Phi_0) = (\psi, \psi, \psi)$. The join irreducibility of Φ and Lemma 5.3 force ψ to be join irreducible.

3. Assume $\Phi \in \mathcal{X}(\mathbf{S}^n, \mathbf{S})$ is join-irreducible, with $\Phi \not\leq h$.

Using Lemma 5.2,3 we get that $(\Phi_0, \Phi_0, \Phi_0 \wedge h)$ and $(0, \Phi_1 \wedge h, \Phi_1)$ are in $\mathcal{X}(\mathbf{S}^n, \mathbf{S})$. Using Lemma 5.1 we get

$$(\Phi_0, \Phi_0, \Phi_0 \wedge h) \vee (0, \Phi_1 \wedge h, \Phi_1) = (\Phi_0, \Phi_0 \vee (\Phi_1 \wedge h), (\Phi_0 \wedge h) \vee \Phi_1) = (\Phi_0, \Phi_h, \Phi_1) = \Phi$$

Therefore, either $\Phi = (\Phi_0, \Phi_0, \Phi_0 \wedge h)$ or $\Phi = (0, \Phi_1 \wedge h, \Phi_1)$. In the first case, take $\psi = \Phi_0$, in the second case, take $\psi = \Phi_1$. Once again, the join irreducibility of Φ and Lemma 5.3, force ψ to be join irreducible. The fact that $\Phi \not\leq h$ yields $\psi \not\leq h$. $\square$

From the previous proposition and Corollary 1, Corollary 2 follows by induction.

Corollary 2. If $\Phi \in \mathcal{X}(\mathbf{S}^n, \mathbf{S})_J$ then $\Phi \wedge h \in \mathcal{X}(\mathbf{S}^n, \mathbf{S})_J$.

Combining the different parts of Proposition 2 we get the following theorem:

Theorem 2. Let $\Phi \in \mathcal{X}(\mathbf{S}^n, \mathbf{S})$.

- (1) Φ is join irreducible if and only if $\Phi = (0, \psi \wedge h, \psi)$ or $\Phi = (\psi, \psi, \psi \wedge h)$ for some join irreducible $\psi \in \mathcal{X}(\mathbf{S}^{n-1}, \mathbf{S})$, i.e. if and only if $\Phi = \Psi \wedge \Pi_1$ or $\Phi = \Psi \wedge \overline{\Pi_1}$, where $\Psi = (\psi, \psi, \psi)$ is the n -ary version of ψ .

- (2) When $\Phi \leq h$, Φ is join irreducible if and only if $\Phi = (0, \psi, \psi)$ or $\Phi = (\psi, \psi, \psi)$ for some join irreducible $\psi \in \mathcal{X}(\underline{\mathbf{S}}^{n-1}, \underline{\mathbf{S}})$ with $\psi \leq h$.
- (3) When $\Phi \not\leq h$, Φ is join irreducible if and only if $\Phi = (0, \psi \wedge h, \psi)$ or $\Phi = (\psi, \psi, \psi \wedge h)$ for some join irreducible $\psi \in \mathcal{X}(\underline{\mathbf{S}}^{n-1}, \underline{\mathbf{S}})$ with $\psi \not\leq h$.

3. THE POSET AND POLYNOMIAL CHARACTERIZATION OF THE JOIN-IRREDUCIBLE MORPHISMS

With Theorem 2 at our disposal, we can now proceed to obtain the poset structure of $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$. We show that this poset structure completely determines $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$ as a partially Stone Space. Along the way we also obtain a polynomial representation.

We are dealing with $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$ with its pointwise partial order inherited from $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})$. This is the same as the open set partial order obtained from the Stone topology. When we discuss the properties of the elements of $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$ covering, being covered or being incomparable, we will be considering them in the poset $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$, not in the partially complemented distributive lattice $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})$.

3.1. The Base of the Hairy Cube.

Proposition 3. *The set $Y^n = \{\Phi \in \mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J \mid \Phi \leq h\}$ is poset isomorphic to 2^n .*

Proof. For $n = 1$ see figure of $\mathcal{X}(\underline{\mathbf{S}}, \underline{\mathbf{S}})_J$ given in Corollary 1.

Assume now that there exists a poset isomorphism $\eta_{n-1} : Y^{n-1} \longrightarrow 2^{n-1}$. By Theorem 2.2 if $\Phi \in Y^n$, then we know that $\Phi = (0, \psi, \psi)$ or $\Phi = (\psi, \psi, \psi)$ for some $\psi \in \mathcal{X}(\underline{\mathbf{S}}^{n-1}, \underline{\mathbf{S}})_J$. Define the following map:

$$\begin{aligned} \eta_n : \quad Y^n &\longrightarrow 2^n \\ (0, \psi, \psi) &\mapsto (0, \eta_{n-1}(\psi)) \\ (\psi, \psi, \psi) &\mapsto (1, \eta_{n-1}(\psi)) \end{aligned}$$

The fact that η_n is bijective follows immediately from the fact that η_{n-1} is. That η_n and its inverse are order preserving is clear from the definition and the fact that η_{n-1} and its inverse are order preserving. $\square$

3.2. The Covers.

Proposition 4. *Let $n \geq 1$ and $\Phi, \Gamma \in \mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$.*

- (1) *If $\Phi, \Gamma \not\leq h$ and $\Phi \neq \Gamma$ then Φ, Γ are incomparable.*
- (2) *If $\Phi \leq h$ then there is a unique $\tilde{\Phi} \in \mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$ with $\tilde{\Phi} \not\leq h$, such that $\tilde{\Phi}$ covers Φ . Moreover $\Phi = \tilde{\Phi} \wedge h$.*
- (3) *If $\Phi \not\leq h$ then it only covers $\Phi \wedge h$ in $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$.*

Proof. (1). The case $n = 1$ is taken care of in Corollary 1. For $n > 1$, Theorem 2.3 tells us that either $\Phi = (0, \phi \wedge h, \phi)$ or $\Phi = (\phi, \phi, \phi \wedge h)$, and similarly $\Gamma = (0, \gamma \wedge h, \gamma)$ or $\Gamma = (\gamma, \gamma, \gamma \wedge h)$ for some $\phi, \gamma \in \mathcal{X}(\underline{\mathbf{S}}^{n-1}, \underline{\mathbf{S}})_J$ with $\phi, \gamma \not\leq h$. Clearly $(0, \phi \wedge h, \phi)$ and $(\gamma, \gamma, \gamma \wedge h)$ are incomparable since $\phi \not\leq h$. By induction $(0, \phi \wedge h, \phi)$ and $(0, \gamma \wedge h, \gamma)$ are

incomparable; similarly, $(\phi, \phi, \phi \wedge h)$ and $(\gamma, \gamma, \gamma \wedge h)$ are incomparable.

(2). Once again, the case $n = 1$ is taken care of in Corollary 1. For $n > 1$, Theorem 2.2 tells us that either $\Phi = (0, \psi, \psi)$ or $\Phi = (\psi, \psi, \psi)$, for some $\psi \in \mathcal{X}(\underline{\mathbf{S}}^{n-1}, \underline{\mathbf{S}})_J$ with $\psi \leq h$. By induction, there is a unique $\tilde{\psi} \in \mathcal{X}(\underline{\mathbf{S}}^{n-1}, \underline{\mathbf{S}})_J$ with $\tilde{\psi} \not\leq h$, such that $\tilde{\psi}$ covers ψ . Moreover, $\psi = \tilde{\psi} \wedge h$. By Theorem 2.3, we have that $(0, \psi, \tilde{\psi})$ and $(\tilde{\psi}, \tilde{\psi}, \psi)$ are in $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$. Clearly, $(0, \psi, \tilde{\psi})$ covers $(0, \psi, \psi)$. That $(\tilde{\psi}, \tilde{\psi}, \psi)$ covers (ψ, ψ, ψ) follows from the fact that neither $(\psi, \tilde{\psi}, \psi)$ nor $(\tilde{\psi}, \psi, \psi)$ are morphisms by Lemma 5.2.d. This shows the existence part, by taking $\tilde{\Phi} = (0, \psi, \tilde{\psi})$ when $\Phi = (0, \psi, \psi)$ and $\tilde{\Phi} = (\tilde{\psi}, \tilde{\psi}, \psi)$ when $\Phi = (\psi, \psi, \psi)$. In either case note that $\Phi = \tilde{\Phi} \wedge h$. For uniqueness, assume that $\Gamma \in \mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$, with $\Gamma \not\leq h$, covers Φ . By Theorem 2.3, we have either $\Gamma = (0, \gamma \wedge h, \gamma)$ or $\Gamma = (\gamma, \gamma, \gamma \wedge h)$ for some $\gamma \in \mathcal{X}(\underline{\mathbf{S}}^{n-1}, \underline{\mathbf{S}})_J$ with $\gamma \not\leq h$. In the first case, it follows that $\Phi = (0, \psi, \psi)$ and γ covers ψ . By uniqueness of $\tilde{\psi}$, we must have $\gamma = \tilde{\psi}$, and $\Gamma = (0, \psi, \tilde{\psi})$. In the second case, since $(0, \psi, \psi) \leq \Gamma$ implies $(\psi, \psi, \psi) \leq \Gamma$ we must have $\Phi = (\psi, \psi, \psi)$ and γ covers ψ . Again, by uniqueness of $\tilde{\psi}$ we get $\gamma = \tilde{\psi}$, and $\Gamma = (\tilde{\psi}, \tilde{\psi}, \psi)$.

(3). By part 1, Φ can only cover elements of $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J \cap \downarrow h$, and part 2 yields the uniqueness. By Corollary 2, $\Phi \wedge h$ is in $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$, and by part 2 it has a unique cover $\tilde{\Phi} \not\leq h$. Therefore, we must have $\tilde{\Phi} \leq \Phi$. Now part 1 forces $\tilde{\Phi} = \Phi$, so Φ covers $\Phi \wedge h$. $\square$

We can now combine the results of Propositions 3 and 4 to describe the poset structure of $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$.

Theorem 3. *The poset $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$ consists of two parts: the “base” Y^n which is an n -cube and the “hairs” $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J \setminus Y^n$, which are pairwise incomparable. Each element of the base is covered by a unique hair. Each hair covers a unique base element.*

In Theorem 4 we show that these partial order properties of $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$ completely determine it as a partially Stone space. Even though we will only need the fact that $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$ is a poset, we can actually see that it is a meet-semilattice.

Corollary 3. *$\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$ is a meet-semilattice. For any $\Phi, \Gamma \in \mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$,*

$$\Phi \wedge \Gamma = (\Phi \wedge h) \wedge (\Gamma \wedge h)$$

is an element of the base of the hairy cube.

3.3. The Partially Stone Space Corresponding to the Hairy Cube.

There is a well-known duality between T_0 Alexandrov spaces and partial orders. For details see [5]. For us, it will be more convenient to use the opposite partial order and the opposite (interchange open and closed) topology.

Here are the details. Given a poset P , the set $\Lambda = \{\downarrow p \mid p \in P\}$ forms a basis for a topology on P ; we refer to it as the “downset topology” (it is the opposite of the “Alexandrov topology”). Given a T_0 -space X , the following defines a partial order on X : for $x, y \in X$, set $x \leq y$ if and only if every open subset of X that contains y must also contain x . We refer to this as the “open set partial order” (it is the opposite of the “specialization order”).

Just like in the Alexandrov duality we get the following lemma:

Lemma 6.

- (1) Suppose that X is a T_0 Alexandrov space, partially ordered with the open set partial order. Then the downset topology induced by the partial order, and the original topology on X are the same.
- (2) Suppose that P is a partially ordered set, which is given the downset topology. Then the open set partial order and the original partial order are the same.

Moreover, a function between posets is order preserving if and only if it is continuous as a map between T_0 Alexandrov spaces.

Note that any Partially Stone Space is T_0 , and every finite space is Alexandrov. Recall that for a finite partially complemented distributive lattice $\mathbb{L}$, the partially Stone space $[X, Y]$ corresponding to $\mathbb{L}$ under the $\mathcal{BSR} \rightleftharpoons \mathcal{PSS}$ duality has $X = \mathbb{L}_J$ and $Y = \{x \in X \mid x \leq h\}$. From the definition of the Stone topology on X and the previous lemma we get the following corollary:

Corollary 4. Let $\mathbb{L}$ be a finite partially complemented distributive lattice. Let $\mathbb{L}_J$ be the set of join-irreducible elements of $\mathbb{L}$, with the partial order inherited from $\mathbb{L}$. Let $[X, Y]$ be the Partially Stone Space corresponding to $\mathbb{L}$ under the duality $\mathcal{BSR} \rightleftharpoons \mathcal{PSS}$.

- (1) The downset topology on $\mathbb{L}_J$ and the Stone topology on X are the same.
- (2) The partial order in $\mathbb{L}_J$ and the open set partial order in X are the same.

We now characterize those partially Stone spaces that correspond to the Hairy Cube.

Theorem 4. Let $[X, Y]$ be a Partially Stone Space partially ordered with the open set partial order such that the following hold:

- (1) Y is poset isomorphic to 2^n ;
- (2) The elements of $X - Y$ are pairwise incomparable.
- (3) Every $y \in Y$ has a unique cover $x \in X - Y$;
- (4) Every $x \in X - Y$ covers only one $y \in Y$;

Then $[X, Y]$, and the Partially Stone Space corresponding to $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})$ under the $\mathcal{BSR} \rightleftharpoons \mathcal{PSS}$ duality, are Partially Stone Space homeomorphic.

Proof. It is clear from Theorem 3 that X and $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$ are poset isomorphic, under an isomorphism $\eta : X \rightarrow \mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$ that maps Y onto Y^n . By Lemma 6, η is a homeomorphism of topological spaces. The only additional fact needed is that $\eta|Y$ is coherent. This follows from the fact that Y and Y^n are finite. $\square$

3.4. Polynomials.

Recall from Lemma 4 that $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$ is just the set of n -ary term functions on the algebra $\mathbb{S}$ which are join-irreducible. The following result shows that these can be obtained using only the $\wedge$ operation and unary $\neg$ operation introduced in Lemma 1.

When writing term functions for an algebra, projection maps are usually called “variables”, and denoted by lower case letters. Given a variable p_i , with $1 \leq i \leq n$, and $\epsilon_i \in \{0, 1\}$ we define

$$p_i^{\epsilon_i} = \begin{cases} p_i & \text{if } \epsilon_i = 0 \\ \overline{p_i} & \text{if } \epsilon_i = 1 \end{cases}$$

Theorem 5. *Let $\Phi \in \mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$.*

(1) *When $\Phi \not\leq h$ it can be uniquely written as a polynomial of the form*

$$\Phi = \bigwedge_{i=1}^n p_i^{\epsilon_i}$$

(2) *When $\Phi \leq h$ it can be uniquely written as a polynomial of the form*

$$\Phi = \left(\bigwedge_{i=1}^n p_i^{\epsilon_i} \right) \wedge h$$

Proof. When $n = 1$ the statement follows from Corollary 1. If $n > 1$ recall the map η in Proposition 3. We will show that taking $\epsilon = \eta(\Phi \wedge h)$, proves existence. By Theorem 2.1 there is $\phi \in \mathcal{X}(\underline{\mathbf{S}}^{n-1}, \underline{\mathbf{S}})_J$, and we have two cases to consider. In the first case, $\Phi = (0, \phi \wedge h, \phi)$,

$$\eta(\Phi \wedge h) = \eta(0, \phi \wedge h, \phi \wedge h) = (0, \eta(\phi \wedge h)), \text{ and } \Phi = (0, h, 1) \wedge (\phi, \phi, \phi) = p_1 \wedge \phi.$$

In the second case, $\Phi = (\phi, \phi, \phi \wedge h)$,


$$\eta(\Phi \wedge h) = \eta(\phi \wedge h, \phi \wedge h, \phi \wedge h) = (1, \eta(\phi \wedge h)), \text{ and } \Phi = (1, 1, h) \wedge (\phi, \phi, \phi) = \overline{p_1} \wedge \phi.$$

In either case, $\Phi \leq h$ if and only if $\phi \leq h$. By induction we have:

$$\text{when } \Phi \not\leq h, \quad \Phi = p_1^{\epsilon_1} \wedge \phi = p_1^{\epsilon_1} \wedge \bigwedge_{i=2}^n p_i^{\eta(\phi \wedge h)} = \bigwedge_{i=1}^n p_i^{\eta(\Phi \wedge h)};$$

$$\text{when } \Phi \leq h, \quad \Phi = p_1^{\epsilon_1} \wedge \phi = p_1^{\epsilon_1} \wedge \bigwedge_{i=2}^n p_i^{\eta(\phi \wedge h)} \wedge h = \bigwedge_{i=1}^n p_i^{\eta(\Phi \wedge h)} \wedge h.$$

Uniqueness follows from the above, the bijectivity of η in Proposition 3, and Proposition 4. $\square$

The “Hairy Cube” for $n = 3$ 

4. NEITHER FULL NOR STRONG, A SMALL STRONG STRUCTURE

From the optimal duality established in Theorem 1, we have obtained geometric and polynomial characterizations of $\mathcal{X}(\underline{\mathbf{S}}^n, \underline{\mathbf{S}})_J$. However, this duality is neither full nor strong, as the following result shows.

Theorem 6. *The duality yielded by $\underline{\mathbf{S}}$ is neither full nor strong.*

Proof. First we will show that the duality yielded by $\underline{\mathbf{S}}$ on $\mathcal{A}$ is not strong. Consider the Second Strong Duality Theorem [1, 3.2.9]. Since $\underline{\mathbf{S}}$ is a total structure, if it were to yield a strong duality on $\mathcal{A}$, it would satisfy the Finite Term Closure Condition:

FTC: For any $n \in \mathbb{N}$, $X \leq \underline{\mathbf{S}}^n$ and $y \in \leq \underline{\mathbf{S}}^n \setminus X$, $\exists$ term functions $\sigma, \tau : \underline{\mathbf{S}}^n \rightarrow \underline{\mathbf{S}}$ on $\underline{\mathbf{S}}$ (that is morphisms) that agree on X but not at y .

Consider $X = \{0, 1\} \leq \underline{\mathbf{S}}$ and $y = h$. Upon viewing the diagram in Lemma 3 we see that $(\underline{\mathbf{S}}, \underline{\mathbf{S}})|_X = \{(0, 0), (0, h), (0, 1), (h, h), (h, 1), (1, h), (1, 1)\}$ and hence no pair $\sigma, \tau \in \mathcal{X}(\underline{\mathbf{S}}, \underline{\mathbf{S}})$ can agree on $\{0, 1\}$ (and differ at h).

To show that the duality yielded on $\mathcal{A}$ by $\underline{\mathbf{S}}$ is not even a full duality, consider the First Strong Duality Theorem [1, 3.2.4], which states that $\underline{\mathbf{S}}$ yields a strong duality on $\mathcal{A}$ if and only if $\underline{\mathbf{S}}$ yields a full duality on $\mathcal{A}$ and $\underline{\mathbf{S}}$ is injective in $\mathcal{X}$. Since $\underline{\mathbf{S}}$ is injective in $\mathcal{X}$, by Theorem 1, it follows that $\underline{\mathbf{S}}$ does not yield a full duality on $\mathcal{A}$. $\square$

The Dual Adjunction Theorem [1, 1.5.3] establishes embeddings of X into $DE(X)$ for every $X \in \mathcal{X}$. The failure of the duality to be full, and therefore strong, is in the failure of X to be isomorphic to $DE(\mathbf{X}) = \mathcal{A}(\mathcal{X}(\mathbf{X}, \underline{\mathbb{S}}), \mathbb{S})$ for every $X \in \mathcal{X}$. In order to obtain a strong duality, we need to add structure to $\underline{\mathbb{S}}$ that will eliminate objects of $\mathcal{X}$ that are a closed substructure of a power of $\underline{\mathbb{S}}$ and are not term/hom-closed.

The NU Strong Duality Theorem, [1, 3.3.8] uses the irreducibility index of $\mathbb{S}$, defined below, to give an exact recipe for constructing a generating structure $\underline{\mathbb{S}}_{\text{NU}}$ that will yield a strong duality on $\mathcal{A}$. One simply needs to add the all the algebraic n -ary operations and partial operations for $1 \leq n \leq \text{Irr}(\mathbb{S})$ to the structure on $\underline{\mathbb{S}}$, to obtain $\underline{\mathbb{S}}_{\text{NU}}$. We refer to this method we refer to as the “Near Brute Force” method. One can then apply the methods of [1], particularly the M-Shift Strong Duality Lemma [1, 3.2.3], to work towards obtaining an optimal strong duality.

Definition 3. *Let A be a finite algebra. The irreducibility of A is the least $n \in \mathbb{N}$ such that the zero congruence on A is a meet of n meet-irreducible congruences. The irreducibility index of A denoted $\text{Irr}(A)$, is the maximum of the irreducibilities of subalgebras of A .*

Lemma 7. *$\text{Irr}(\mathbb{S})$ is 2.*

Proof. From the lattice of subalgebras of $\mathbb{S}^2$ in Lemma 2 it is easy to check that $\text{Con}(\mathbb{S}) = \{\Delta, r_3, r_2 \cap r_2^{-1}, \mathbb{S}^2\}$, and it is isomorphic to the 2 dimensional cube. $\mathbb{S}$ has no subalgebra other than itself. $\square$

Recall that a n -ary operation g on S is algebraic over $\mathbb{S}$ if $g \in \mathcal{A}(\mathbb{S}^n, \mathbb{S})$; a n -ary partial operation h on S is algebraic over $\mathbb{S}$ if $h \in \mathcal{A}(X, \mathbb{S})$ for some $X \leq \mathbb{S}^n$. Furthermore, these conditions are equivalent to saying that the corresponding graphs form subalgebras of $\mathbb{S}^{n+1}$. As can be seen through the proofs of Lemma 8 and Proposition 6, the number of algebraic binary partial operations on $\mathbb{S}$ is too large, for the brute force method to yield a useful structure. We want a structure $\underline{\mathbb{S}}_{\text{s}}$ that yields a strong duality on $\mathcal{A}$ that is as simple as possible. On the other hand, the only algebraic binary total operations on $\mathbb{S}$ are the projections.

Proposition 5. *Let $n \in \mathbb{N}$ and $\Lambda : \mathbb{S}^n \rightarrow \mathbb{S}$. Then $\Lambda \in \mathcal{A}(\mathbb{S}^n, \mathbb{S})$ if and only if it is a projection map, $\Lambda = \Pi_i^n$, for some $1 \leq i \leq n$.*

Proof. $\Lambda^{-1}(1)$ (resp. $\Lambda^{-1}(0)$) is a prime filter (resp. ideal) of $\mathbb{S}^n$, hence there is x (resp. y) join-irreducible (resp. meet-irreducible) such that $\Lambda^{-1}(1) = \uparrow x$ (resp. $\Lambda^{-1}(0) = \downarrow y$). As x is join-irreducible, it follows that $\Pi_j(x) \neq 0$ for at most one j . Since h is a constant, $\Lambda(x \wedge h) = \Lambda(x) \wedge h = 1 \wedge h = h$, and we cannot have $x \wedge h = x$ and therefore $\Pi_j(x) = 1$. Similarly, $\Pi_i(y) \neq 1$ for at most one i , and $\Pi_i(y) = 0$. If $i \neq j$, then $x \leq y$ and $\Lambda(x) \leq \Lambda(y)$ yielding a contradiction, hence $i = j$. Now, let $z \in X$ and consider the following cases:

- (1) $\Pi_i(z) = 0$. In this case $z \in \downarrow y$ and hence $\Phi(z) = 0$.
- (2) $\Pi_i(z) = h$. In this case $z \notin \uparrow x$ and $z \notin \downarrow y$, hence $\Phi(z) = h$.
- (3) $\Pi_i(z) = 1$. In this case $z \in \uparrow x$ and hence $\Phi(z) = 1$.

So we see that $\Phi = \Pi_i$. The converse holds as projections are homomorphisms. $\square$

As a result of Proposition 5 the only total operations in $\underline{\mathbf{S}}_{\text{NU}}$ are Π_1^1 , Π_1^2 and Π_2^2 . There are no proper algebraic unary partial operations, since the only subalgebra of $\mathbf{S}$ is $\mathbf{S}$ itself, but the set of algebraic binary partial operations is too large to be useful.

In order to get a manageable structure $\underline{\mathbf{S}}_{\text{s}}$ that yields a strong duality on $\mathcal{A}$, we will reduce the set of algebraic binary partial operations using the M-Shift Strong Duality Lemma. Any structure that strongly entails $\underline{\mathbf{S}}_{\text{NU}}$ will also yield a strong duality on $\mathcal{A}$. To get such structure $\underline{\mathbf{S}}_{\text{s}}$, we may delete from $\underline{\mathbf{S}}_{\text{NU}}$ those partial operations that are restrictions of other total or partial operations left in the structure. In particular, we may delete any partial operation which is a restriction of a projection.

Unlike total algebraic operations, which by Proposition 5 have to be projections, for the algebraic binary partial operations there is a little more room as the following lemma shows.

Lemma 8. *Let $A \leq \mathbb{S}^2$. For $i = 1, 2$, let $L_i = \Pi_i^2$ restricted to $\downarrow h$, $U_i = \Pi_i^2$ restricted to $\uparrow h$. Let $\Lambda \in \mathcal{A}(A, \mathbf{S})$.*

- (1) *If $A \cap \uparrow h$ contains $(h, 1)$ or $(1, h)$ then $\uparrow h \subseteq A$ and Λ restricted to $\uparrow h$ equals U_1 or U_2 .*
- (2) *Λ restricted to $A \cap \downarrow h$ equals L_1 or L_2 .*

Proof. If $A \cap \uparrow h$ contains either $(h, 1)$ or $(1, h)$, applying the complement operation from Lemma 1 we get the other and hence $\uparrow h \subseteq A$. The rest follows from the facts that Λ is order preserving and every element of $\Delta_{\mathbf{S}}$ is a constant. $\square$

It is easy to check that the binary partial operation λ_1 with domain r_1 having graph

$$\Gamma(\lambda_1) = \{(0, 0, 0), (h, h, h), (1, 1, 1), (0, h, h), (0, 1, h), (h, 0, 0), (h, 1, h), (1, h, 1)\},$$

is in fact algebraic. It combines U_1 and L_2 . Similarly, the combination of U_2 and L_1 yields the algebraic binary partial operation λ_2 with domain r_1^{-1} having graph

$$\Gamma(\lambda_2) = \{(0, 0, 0), (h, h, h), (1, 1, 1), (0, h, 0), (h, 0, h), (1, 0, h), (h, 1, 1), (1, h, h)\}.$$

These two partial operations are examples of algebraic binary partial operations which are not restrictions of projections. Lemma 8 places constraints on such operations to be combinations of L_i and U_k with $i \neq k$. In a sense, λ_1 and λ_2 are the only such examples, as it is more clearly stated in the proof of the next proposition.

Proposition 6. *Let $n \in \mathbb{N}$ and $\Pi_i^n : \underline{\mathbf{S}}^n \rightarrow \underline{\mathbf{S}}$ denote the i -th projection map for any $i \in \mathbb{N}$ with $1 \leq i \leq n$ and $\mathcal{T}$ denote the discrete topology, then the structure*

$$\underline{\mathbf{S}}_{\text{s}} = \langle S; \{r_1, r_2, r_3\}, \{\Pi_1^2, \Pi_2^2\}, \{\lambda_1, \lambda_2\}, \mathcal{T} \rangle$$

yields a strong duality on $\mathcal{A}$.

Proof. As Π_1^1 is the identity map on S , it has no effect on any topological category generated by a structure with S as its carrier set. Therefore, we do not need to include it in the list of total operations. By the M-Shift Strong Duality Lemma we only need to show that any algebraic binary partial operation in the structure $\underline{\mathbf{S}}_{\text{NU}}$ is a restriction of either a projection or one of λ_1, λ_2 . From Lemma 8, we see that the only homomorphisms which are not restrictions of projections must consist of either a combination of L_1 and U_2 or a combination of L_2 and

U_1 . Let us first consider the subalgebra $A = r_1 \leq \mathbb{S}^2$ which contains the element $(0, 1)$. Let $\lambda : A \rightarrow \mathbb{S}$ be a homomorphism which is not a restriction of a projection. If we had $\lambda(0, 1) = 0$ this would force $\lambda(0, h) = \lambda(h, h) \wedge \lambda(0, 1) = h \wedge 0 = 0$ and by Lemma 8, λ restricted to $A \cap \downarrow h$ would have to equal L_1 . Moreover, we would have $\lambda(h, 1) = \lambda(h, h) \vee \lambda(0, 1) = h \vee 0 = h$, and λ restricted to $A \cap \uparrow h$ would have to equal U_1 , making λ a restriction of Π_1^2 . Similarly, if we had $\lambda(0, 1) = 1$ this would force λ to be a restriction of Π_2^2 . Therefore, we must have $\lambda(0, 1) = h$. As above, this forces $\lambda(0, h) = \lambda(h, h) \wedge \lambda(0, 1) = h \wedge h = h$, and $\lambda(h, 1) = \lambda(h, h) \vee \lambda(0, 1) = h \vee h = h$, making λ a combination of L_2 and U_1 , i.e. λ_1 . So, λ_1 is the only partial algebraic operation with domain r_1 which is not a restriction of a projection. A similar argument shows that λ_2 is the only partial algebraic operation with domain r_1^{-1} , which is not a restriction of a projection.

Note that the argument above does not make use of the fact that $(h, 0)$ is in r_1 . Therefore, it also shows that the only partial algebraic operation with domain r_2 which is not a restriction of a projection, must be the restriction of λ_1 . Similarly, the only partial algebraic operation with domain r_2^{-1} which is not a restriction of a projection, must be the restriction of λ_2 .

As shown in Lemma 2 any other subalgebra $A \leq \mathbb{S}^2$ must be a subalgebra of $r_1 \cap r_1^{-1}$. Hence, by Lemma 8, any partial algebraic operation with domain A , which is not a restriction of a projection, must be a restriction of either λ_1 or λ_2 . $\square$

As an intermediate step towards an optimal strong duality, we will show that we can eliminate the total operations and λ_2 from $\underline{\mathbf{S}}_s$ and still achieve a strong duality. First we need the following definitions:

Definition 4. [1] *A set of partial operations on a set X is called an enriched partial clone if it is closed under composition and contains $\{\Pi_i^n \mid 1 \leq i \leq n, n \in \mathbb{N}\}$. The enriched partial clone generated by a set of partial operations H on a set X is the smallest enriched partial clone containing H and is denoted $[H]$. Given a structured topological space $\mathbf{X} = \langle X; R, G, H, T \rangle$, the enriched partial clone of $\mathbf{X}$ is the smallest enriched partial clone on X containing $G \cup H$ and is denoted $[G \cup H]$. Furthermore, let $\mathcal{P}$ denote the set of all finitary algebraic, partial or total operations on $\mathbb{S}$. Then $\mathcal{P}$ is an enriched partial clone on S and is referred to as the enriched partial hom-clone of $\mathbb{S}$.*

Definition 5. [1] *Let $P \subseteq \mathcal{P}$ and $k \in \mathcal{P}$. We say that P hom-entails k if, for all non-empty sets Ω , each topologically closed subset of S^Ω which is closed under the partial operations in P is also closed under k . Define $\overline{P} = \{k \in \mathcal{P} \mid P \text{ hom-entails } k\}$. Then $P \mapsto \overline{P}$ is a closure operator on $\mathcal{P}$ and we refer to $\overline{P}$ as the hom-entailment closure of P .*

Corollary 5. *The structure $\underline{\mathbf{S}}'_s = \langle S; \{r_1, r_2, r_3\}, \{\lambda_1\}, T \rangle$ yields a strong duality on $\mathcal{A}$.*

Proof. Let $G = \{\Pi_1^2, \Pi_2^2\}$ and $H = \{\lambda_1, \lambda_2\}$, then by Lemma 6 and the Brute Force Strong Duality Theorem [1, 3.2.2], $\overline{G \cup H} = \mathcal{P}$, i.e. $G \cup H$ hom-entails every finitary algebraic partial or total operation on $\mathbb{S}$. Now consider Definition 4, and note that $[H] = [G \cup H]$. Further note that $\lambda_1(\Pi_2^2, \Pi_1^2) = \lambda_2$, and we see that $[\lambda_1] = [H]$. Let $k : A \rightarrow S$ be any element of $\mathcal{P}$ and consider what we refer to as the Test Algebra Lemma for maps [1, 9.4.1]. Parts (i.a) and (i.c) show $\overline{\{\lambda_1\}} = \mathcal{P}$ and hence the structure $\underline{\mathbf{S}}'_s$ yields a strong duality on $\mathcal{A}$. $\square$

Now we wish to show that the partial operation λ_1 entails r_1 and r_3 , for that we need the following Lemma:

Lemma 9. *Let $\underline{\mathbf{S}}' = \langle S; \{\lambda_1\}, \mathcal{T} \rangle$ with λ_1 as the only partial operation.*

- (1) *Let $\mathbf{X} \in IS_cP^+(\underline{\mathbf{S}}')$ and $g : \mathbf{X} \longrightarrow \underline{\mathbf{S}}'$ be a morphism. Then g preserves r_1 and r_3 , and therefore $\{\lambda_1\}$ entails r_1 and r_3 .*
- (2) *Now, let $\Phi : \underline{\mathbf{S}}' \longrightarrow \underline{\mathbf{S}}'$ be given by $\Phi = (h, 0, 0)$. Then Φ preserves λ_1 but not r_2 , and therefore $\underline{\mathbf{S}}'$ does not yield a duality on $\mathcal{A}$.*

Proof. (1) Let $(Y, Z) \in r_1^{\mathbf{X}} = \text{dom}(\lambda_1^{\mathbf{X}})$. As g preserves λ_1 , $(g(Y), g(Z)) \in \text{dom}(\lambda_1^{\underline{\mathbf{S}}'}) = r_1^{\underline{\mathbf{S}}'}$ and g preserves r_1 . Now let $(Y, Z) \in r_3^{\mathbf{X}} \subseteq \text{dom}(\lambda_1^{\mathbf{X}})$, and consider $(Y, Z, \lambda_1(Y, Z))$. For any i , $(Y_i, Z_i) \in r_3^{\underline{\mathbf{S}}'}$ hence $Z_i = \lambda_1(Y_i, Z_i)$ and therefore $\lambda_1(Y, Z) = Z \in \mathbf{X}$. Moreover, $g(Z) = g(\lambda_1(Y, Z))$ which implies that $(g(Y), g(Z)) \in r_3^{\underline{\mathbf{S}}'}$. The fact that $\{\lambda_1\}$ entails r_1 and r_3 follows from the Entailment Lemma.

(2) The fact that Φ preserves λ_1 is easily seen by inspection of $\Phi(\lambda_1(x, y))$ and $\lambda_1(\Phi(x), \Phi(y))$ for each $(x, y) \in \text{dom}(\lambda_1)$. Now note that $(\Phi(0), \Phi(h)) = (h, 0)$ and hence Φ fails to preserve r_2 . From Lemmas 3 and 4 we see that $\underline{\mathbf{S}}'$ does not yield duality on $\mathcal{A}$. $\square$

We now have enough results to prove the major Theorem of this section.

Theorem 7. *Let $\mathcal{T}$ denote the discrete topology and $\underline{\mathbf{S}}_{os} = \langle S; \{r_2\}, \{\lambda_1\}, \mathcal{T} \rangle$, then $\underline{\mathbf{S}}_{os}$ yields an optimal strong duality on $\mathcal{A}$.*

Proof. The fact that $\underline{\mathbf{S}}_{os}$ yields a strong duality on $\mathcal{A}$ follows from Theorem 1, Corollary 5, Lemma 9.1 and the M-Shift Strong Duality Lemma. The fact that it is optimal follows from Lemma 9.2 and Theorem 1. $\square$

Combining this theorem with Lemma 4 we get the final result.

Corollary 6. *Let $\mathcal{X}_{os} = IS_cP^+(\underline{\mathbf{S}}_{os})$, then for any $n \in \mathbb{N}$, $\mathcal{X}_{os}(\underline{\mathbf{S}}_{os}^n, \underline{\mathbf{S}})_J$ is the n -dimensional Hairy Cube.*

REFERENCES

References

- [1] David M. Clark and Brian A. Davey, *Natural dualities for the working algebraist*, Cambridge Studies in Advanced Mathematics, vol. 57, Cambridge University Press, Cambridge, 1998. MR **1663208** (2000d:18001)
- [2] Daniel J. Clouse, *A Dual Representation of Boolean Semirings in a Category of Structured Topological Spaces*, Ph.D. dissertation, Binghamton University, 2002.
- [3] Samuel Eilenberg, *Automata, languages, and machines. Vol. A*, Academic Press [A subsidiary of Harcourt Brace Jovanovich, Publishers], New York, 1974. Pure and Applied Mathematics, Vol. 58. MR 0530382 (58 #26604a)
- [4] Fernando Guzmán, *The variety of Boolean semirings*, J. Pure Appl. Algebra **78** (1992), no. 3, 253–270. MR **1163278** (93d:08007)
- [5] Peter T. Johnstone, *Stone spaces*, Cambridge Studies in Advanced Mathematics, vol. 3, Cambridge University Press, Cambridge, 1982. MR **698074** (85f:54002)

- [6] Werner Kuich and Arto Salomaa, *Semirings, automata, languages*, EATCS Monographs on Theoretical Computer Science, vol. 5, Springer-Verlag, Berlin, 1986. MR **817983** (**87h**:68093)
- [7] M. H. Stone, *Applications of the theory of Boolean rings to general topology*, Trans. Amer. Math. Soc. **41** (1937), no. 3, 375–481. MR 1501905

DANIEL J. CLOUSE

DEPARTMENT OF DEFENSE

E-mail address: `beckclouse@netzero.net`

FERNANDO GUZMÁN

DEPT. OF MATHEMATICAL SCIENCES, BINGHAMTON UNIVERSITY, BINGHAMTON N.Y., 13902-6000

E-mail address: `fer@math.binghamton.edu`

URL: `http://math.binghamton.edu/fer`